

## Do I need to see a doctor or a therapist?

Not usually – if you follow the right advice and take the right medication, your back problem should improve over the next 6 weeks. If your back symptoms are still interfering with your ability to do normal things after 6 weeks, this is the time to seek further professional advice from either your GP or NHS Physiotherapist.

## What should I look out for?

You should return to your local Emergency Department as soon as possible if you develop any of these:

- Difficulty passing or controlling urine
- Numbness and/or altered sensation such as pins and needles around your back passage or genitals, e.g. when wiping after toileting.

If you experience any of the following in conjunction with your back pain, you need to speak to your doctor as soon as possible:

- Generally feeling unwell
- Back pain that starts when you have other problems, such as rheumatoid arthritis or cancer

- Numbness, pins and needles, or weakness in one or both legs that has not improved after one week
- Unsteadiness when you walk
- Your back problem has not improved within six weeks.

## Facts and figures

- 90% of the UK population get back pain at some point
- Most back pain settles within six weeks
- Keeping active is the best thing to help your back pain.

## Further information

- <http://www.noebackpainprogramme.nhs.uk/>
- NHS Inform <http://m.nhsinform.co.uk/health-library/articles/b/back-pain/introduction>
- The truth about back pain - AXA PPP [www.youtube.com/watch?v=b-cBtPSf0Hc](http://www.youtube.com/watch?v=b-cBtPSf0Hc)
- NHS24 MSK help: useful exercises and advice - App for iPhone, Android

© G16041503W. Design Services  
Salford Royal NHS Foundation Trust  
All Rights Reserved 2016  
This document **MUST NOT** be photocopied  
Information Leaflet Control Policy:  
Unique Identifier: MED 23 (16)  
Review Date: May 2018

Salford Royal **NHS**  
NHS Foundation Trust  
University Teaching Trust

safe • clean • personal

For further information on this leaflet, it's references and sources used, please contact **0161 206 4841**

Copies of this information are available in other languages and formats upon request. In accordance with the Equality Act we will make 'reasonable adjustments' to enable individuals with disabilities, to access this treatment / service.

### If you need this interpreting please telephone

Polish  
Jeżeli potrzebne jest Państwu to tłumaczenie, proszę zadzwonić pod numer.

Urdu  
اگر آپ کو اس ترجمانی کی ضرورت ہے تو مہربانی کر کے فون کریں۔

Arabic  
إذا كنتم بحاجة الى تفسير او ترجمة هذا الرجاء الاتصال

Chinese  
如果需要翻译，请拨打电话

Farsi  
اگر به ترجمه این نیاز دارید ، لطفاً تلفن کنید

**0161 206 0224**

Email: [InterpretationandTrans@srft.nhs.uk](mailto:InterpretationandTrans@srft.nhs.uk)

### Salford Royal operates a smoke-free policy.

For advice on stopping smoking contact the Hospital Specialist Stop Smoking Service on 0161 206 1779

Salford Royal NHS Foundation Trust  
Stott Lane,  
Salford,  
Manchester,  
M6 8HD

Telephone  
**0161 789 7373**  
[www.srft.nhs.uk](http://www.srft.nhs.uk)

If you would like to become a Foundation Trust Member please visit:

[www.srft.nhs.uk/for-members](http://www.srft.nhs.uk/for-members)

If you have any suggestions as to how this document could be improved in the future then please visit:

<http://www.srft.nhs.uk/for-patients>

# Back Pain



Hope Building  
Emergency Medicine  
**0161 206 4841**

Health & care  
information  
you can trust  
The Information Standard Certified Member

Salford Royal **NHS**  
NHS Foundation Trust  
University Teaching Trust

safe • clean • personal

## Back Pain

Back problems are very common. Most back problems start for no obvious reason, which can be very frustrating. Your back problem may cause aching, hot, burning, shooting, or stabbing pains in your back and sometimes into one or both of your legs. You may also get pins and needles. The spine is strong and back problems are rarely due to any serious disease or damage. Back problems should settle within 6 weeks of following the advice provided here.

**You will not normally need an X-ray or an MRI scan.**



## What should I do?

- Keep moving, even if slowly at first
- Keep living and working normally. This is important and is the best way to get better
- Don't worry if your back still hurts at work, consider doing light tasks at first, speak to your manager about work concerns that you may have
- Don't sit down for too long, change positions regularly wherever you are
- Avoid bed rest during the day
- Stay active and remember to re-introduce activities like heavy lifting gradually
- Exercise really helps your back and can relieve pain, start with light fitness training. Moving will make you stronger, keeping active is the best thing you can do.

## What about pain relief?

- Initially it may be helpful to use a covered icepack to ease your pain - never apply ice directly onto your skin. Alternatively heat can be soothing, so a covered hot water bottle may also be used. You should not use heat /ice for more than 15 minutes, three to four times a day
- Painkillers may help you keep moving, so sensible use of painkillers such as paracetamol, and ibuprofen will help, not harm your back. However, if you are already taking medication for something else or have other health problems, check with your local pharmacist before taking painkillers. Always follow the instructions on the packet.
- Occasionally, stronger pain-killers such as codeine may be recommended. If you are taking these, you should drink plenty of fluids and eat foods high in fibre (such as fruit and vegetables) to avoid constipation.
- Diazepam or similar drugs are occasionally prescribed for 1-2 days to relax the muscles of the lower back.

## What about work, sports?

You will recover faster if you can stay at or get back to work as early as possible. Don't worry if your back still hurts; consider doing light tasks at first if this helps you get back to work easier and quicker. Try to stay active and remember to keep moving. Speak to your manager at work about any concerns you may have. You should try to do your normal activities as much as possible and use painkillers as needed. With regard to sports, start with light fitness training, and play when you feel ready.

## Can my back problem cause trouble anywhere else?

Sometimes when you get back pain you can also get pain in one or both legs (sciatica). This can cause pins and needles or a shooting, stabbing pain to the legs. This can be very painful but this is common when you have back pain.